

ON HYPERCOMPLEX PSEUDO-HERMITIAN MANIFOLDS

KOSTADIN GRIBACHEV

*Faculty of Mathematics and Informatics, University of Plovdiv,
236 Bulgaria Blvd., Plovdiv 4003, Bulgaria
E-mail: costas@uni-plovdiv.bg*

MANCHO MANEV

*Faculty of Mathematics and Informatics, University of Plovdiv,
236 Bulgaria Blvd., Plovdiv 4003, Bulgaria
E-mail: mmanev@uni-plovdiv.bg*

STANCHO DIMIEV

*Institute of Mathematics and Informatics, Bulgarian Academy of Sciences,
Acad. G. Bonchev Str., Bl. 8, Sofia 1113, Bulgaria,
E-mail: sdimiev@math.bas.bg*

The class of the hypercomplex pseudo-Hermitian manifolds is considered. The flatness of the considered manifolds with the 3 parallel complex structures is proved. Conformal transformations of the metrics are introduced. The conformal invariance and the conformal equivalence of the basic types manifolds are studied. A known example is characterized in relation to the obtained results.

Introduction

This paper is a continuation of the same authors's paper ⁴ which is inspired by the seminal work ¹ of D. V. Alekseevsky and S. Marchiafava. We follow a parallel direction including skew-Hermitian metrics with respect to the almost hypercomplex structure.

In the first section we give some necessary facts concerning the almost hypercomplex pseudo-Hermitian manifolds introduced in ⁴.

In the second one we consider the special class of (integrable) hypercomplex pseudo-Hermitian manifolds, namely pseudo-hyper-Kähler manifolds. Here we expose the proof of the mentioned in ⁴ statement that each pseudo-hyper-Kähler manifold is flat.

The third section is fundamental for this work. A study of the group of conformal transformations of the metric is initiated here. The conformal invariant classes and the conformal equivalent class to the class of the pseudo-hyper-Kähler manifolds are found.

Finally, we characterize a known example in terms of the conformal transformations.

1 Preliminaries

1.1 Hypercomplex pseudo-Hermitian structures in a real vector space

Let V be a real $4n$ -dimensional vector space. By $\left\{ \frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial u^i}, \frac{\partial}{\partial v^i} \right\}$, $i = 1, 2, \dots, n$, is denoted a (local) basis on V . Each vector x of V is represented in the mentioned basis as follows

$$x = x^i \frac{\partial}{\partial x^i} + y^i \frac{\partial}{\partial y^i} + u^i \frac{\partial}{\partial u^i} + v^i \frac{\partial}{\partial v^i}. \quad (1)$$

A standard complex structure on V is defined as in ⁵:

$$\begin{aligned} J_1 \frac{\partial}{\partial x^i} &= \frac{\partial}{\partial y^i}, & J_1 \frac{\partial}{\partial y^i} &= -\frac{\partial}{\partial x^i}, & J_1 \frac{\partial}{\partial u^i} &= -\frac{\partial}{\partial v^i}, & J_1 \frac{\partial}{\partial v^i} &= \frac{\partial}{\partial u^i}; \\ J_2 \frac{\partial}{\partial x^i} &= \frac{\partial}{\partial u^i}, & J_2 \frac{\partial}{\partial y^i} &= \frac{\partial}{\partial v^i}, & J_2 \frac{\partial}{\partial u^i} &= -\frac{\partial}{\partial x^i}, & J_2 \frac{\partial}{\partial v^i} &= -\frac{\partial}{\partial y^i}; \\ J_3 \frac{\partial}{\partial x^i} &= -\frac{\partial}{\partial v^i}, & J_3 \frac{\partial}{\partial y^i} &= \frac{\partial}{\partial u^i}, & J_3 \frac{\partial}{\partial u^i} &= -\frac{\partial}{\partial y^i}, & J_3 \frac{\partial}{\partial v^i} &= \frac{\partial}{\partial x^i}. \end{aligned} \quad (2)$$

The following properties about J_α are direct consequences of (2)

$$\begin{aligned} J_1^2 &= J_2^2 = J_3^2 = -Id, \\ J_1 J_2 &= -J_2 J_1 = J_3, & J_2 J_3 &= -J_3 J_2 = J_1, & J_3 J_1 &= -J_1 J_3 = J_2. \end{aligned} \quad (3)$$

If $x \in V$, i.e. $x(x^i, y^i, u^i, v^i)$ then according to (2) and (3) we have

$$J_1 x(-y^i, x^i, v^i, -u^i), \quad J_2 x(-u^i, -v^i, x^i, y^i), \quad J_3 x(v^i, -u^i, y^i, -x^i). \quad (4)$$

Definition 1.1 (1) A triple $H = (J_1, J_2, J_3)$ of anticommuting complex structures on V with $J_3 = J_1 J_2$ is called a hypercomplex structure on V ;

A bilinear form f on V is defined as ordinary, $f : V \times V \rightarrow \mathbb{R}$. We denote by $\mathcal{B}(V)$ the set of all bilinear forms on V . Each f is a tensor of type $(0, 2)$, and $\mathcal{B}(V)$ is a vector space of dimension $16n^2$.

Let J be a given complex structure on V . A bilinear form f on V is called *Hermitian* (respectively, *skew-Hermitian*) with respect to J if the identity $f(Jx, Jy) = f(x, y)$ (respectively, $f(Jx, Jy) = -f(x, y)$) holds true.

Definition 1.2 (1) A bilinear form f on V is called an Hermitian bilinear form with respect to $H = (J_\alpha)$ if it is Hermitian with respect to any complex structure J_α , $\alpha = 1, 2, 3$, i.e.

$$f(J_\alpha x, J_\alpha y) = f(x, y) \quad \forall x, y \in V. \quad (5)$$

We denote by $L_0 = \mathcal{B}_H(V)$ the set of all Hermitian bilinear forms on V . The notion of pseudo-Hermitian bilinear forms is introduced by the following

Definition 1.3 ⁽⁴⁾ A bilinear form f on V is called a pseudo-Hermitian bilinear form with respect to $H = (J_1, J_2, J_3)$, if it is Hermitian with respect to J_α and skew-Hermitian with respect to J_β and J_γ , i.e.

$$f(J_\alpha x, J_\alpha y) = -f(J_\beta x, J_\beta y) = -f(J_\gamma x, J_\gamma y) = f(x, y) \quad \forall x, y \in V, \quad (6)$$

where (α, β, γ) is a circular permutation of $(1, 2, 3)$.

We denote $f \in L_\alpha \subset \mathcal{B}(V)$ ($\alpha = 0, 1, 2, 3$) when f satisfies the conditions (5) and (6), respectively.

In ¹ is introduced a pseudo-Euclidian metric g with signature $(2n, 2n)$ as follows

$$g(x, y) := \sum_{i=1}^n (-x^i a^i - y^i b^i + u^i c^i + v^i d^i), \quad (7)$$

where $x(x^i, y^i, u^i, v^i)$, $y(a^i, b^i, c^i, d^i) \in V$, $i = 1, 2, \dots, n$. This metric satisfies the following properties

$$g(J_1 x, J_1 y) = -g(J_2 x, J_2 y) = -g(J_3 x, J_3 y) = g(x, y). \quad (8)$$

This means that the pseudo-Euclidean metric g belongs to L_1 .

The form $g_1 : g_1(x, y) = g(J_1 x, y)$ coincides with the Kähler form Φ which is Hermitian with respect to J_α , i.e.

$$\Phi(J_\alpha x, J_\alpha y) = \Phi(x, y), \quad \alpha = 1, 2, 3, \quad \Phi \in L_0.$$

The attached to g associated bilinear forms $g_2 : g_2(x, y) = g(J_2 x, y)$ and $g_3 : g_3(x, y) = g(J_3 x, y)$ are symmetric forms with the properties

$$\begin{aligned} -g_2(J_1 x, J_1 y) &= -g_2(J_2 x, J_2 y) = g_2(J_3 x, J_3 y) = g_2(x, y), \\ -g_3(J_1 x, J_1 y) &= g_3(J_2 x, J_2 y) = -g_3(J_3 x, J_3 y) = g_3(x, y), \end{aligned} \quad (9)$$

i.e. $g_2 \in L_3$, $g_3 \in L_2$.

It follows that the Kähler form Φ is Hermitian regarding H and the metrics g, g_2, g_3 are pseudo-Hermitian of different types with signature $(2n, 2n)$.

Now we recall the following notion:

Definition 1.4 ⁽⁴⁾ The structure $(H, G) := (J_1, J_2, J_3, g, \Phi, g_2, g_3)$ is called a hypercomplex pseudo-Hermitian structure on V .

1.2 Structural tensors on an almost (H, G) -manifold

Let (M, H) be an almost hypercomplex manifold ¹. We suppose that g is a symmetric tensor field of type $(0, 2)$. If it induces a pseudo-Hermitian inner product in $T_p M$, $p \in M$, then g is called a pseudo-Hermitian metric on M .

The structure $(H, G) := (J_1, J_2, J_3, g, \Phi, g_2, g_3)$ is called an *almost hypercomplex pseudo-Hermitian structure on M* or in short an *almost (H, G) -structure on M* . The manifold M equipped with H and G , i.e. (M, H, G) , is called an *almost hypercomplex pseudo-Hermitian manifold*, or in short an *almost (H, G) -manifold*.⁴

The 3 tensors of type $(0, 3)$ $F_\alpha : F_\alpha(x, y, z) = g((\nabla_x J_\alpha)y, z)$, $\alpha = 1, 2, 3$, where ∇ is the Levi-Civita connection generated by g , is called *structural tensors of the almost (H, G) -manifold*.⁴

The structural tensors satisfy the following properties:

$$\begin{aligned} F_1(x, y, z) &= F_2(x, J_3y, z) + F_3(x, y, J_2z), \\ F_2(x, y, z) &= F_3(x, J_1y, z) + F_1(x, y, J_3z), \\ F_3(x, y, z) &= F_1(x, J_2y, z) - F_2(x, y, J_1z); \end{aligned} \quad (10)$$

$$\begin{aligned} F_1(x, y, z) &= -F_1(x, z, y) = -F_1(x, J_1y, J_1z), \\ F_2(x, y, z) &= F_2(x, z, y) = F_2(x, J_2y, J_2z), \\ F_3(x, y, z) &= F_3(x, z, y) = F_3(x, J_3y, J_3z). \end{aligned} \quad (11)$$

Let us recall the Nijenhuis tensors $N_\alpha(X, Y) = \frac{1}{2} [[J_\alpha, J_\alpha]](X, Y)$ for almost complex structures J_α and $X, Y \in \mathfrak{X}(M)$, where

$$[[J_\alpha, J_\alpha]](X, Y) = 2\{[J_\alpha X, J_\alpha Y] - J_\alpha[J_\alpha X, Y] - J_\alpha[X, J_\alpha Y] - [X, Y]\}.$$

It is well known that the almost hypercomplex structure $H = (J_\alpha)$ is a hypercomplex structure if $[[J_\alpha, J_\alpha]]$ vanishes for each $\alpha = 1, 2, 3$. Moreover it is known that one almost hypercomplex structure H is hypercomplex if and only if two of the structures J_α ($\alpha = 1, 2, 3$) are integrable. This means that two of the tensors N_α vanish.¹

We recall also the following definitions. Since g is Hermitian metric with respect to J_1 , according to³ the class $\mathcal{W}_4$ is a subclass of the class of Hermitian manifolds. If (H, G) -manifold belongs to $\mathcal{W}_4$, with respect to J_1 , then the almost complex structure J_1 is integrable and

$$F_1(x, y, z) = \frac{1}{2(2n-1)} [g(x, y)\theta_1(z) - g(x, z)\theta_1(y) - g(x, J_1y)\theta_1(J_1z) + g(x, J_1z)\theta_1(J_1y)], \quad (12)$$

where $\theta_1(\cdot) = g^{ij}F_1(e_i, e_j, \cdot) = \delta\Phi(\cdot)$ for the basis $\{e_i\}_{i=1}^{4n}$, and δ – the coderivative.

On other side the metric g is a skew-Hermitian with respect to J_2 and J_3 , i.e. $g(J_2x, J_2y) = g(J_3x, J_3y) = -g(x, y)$. A classification of all almost complex manifolds with skew-Hermitian metric (Norden metric or B-metric) is given in². One of the basic classes of integrable almost complex manifolds

with skew-Hermitian metric is $\mathcal{W}_1$. It is known that if an almost (H, G) -manifold belongs to $\mathcal{W}_1(J_\alpha)$, $\alpha = 2, 3$, then J_α is integrable and the following equality holds

$$F_\alpha(x, y, z) = \frac{1}{4n} [g(x, y)\theta_\alpha(z) + g(x, z)\theta_\alpha(y) + g(x, J_\alpha y)\theta_\alpha(J_\alpha z) + g(x, J_\alpha z)\theta_\alpha(J_\alpha y)], \quad (13)$$

where $\theta_\alpha(z) = g^{ij}F_\alpha(e_i, e_j, z)$, $\alpha = 2, 3$, for an arbitrary basis $\{e_i\}_{i=1}^{4n}$.

When (12) is satisfied for (M, H, G) , we say that $(M, H, G) \in \mathcal{W}(J_1)$. In the case, (M, H, G) satisfies (13) for $\alpha = 2$ or $\alpha = 3$, we say $(M, H, G) \in \mathcal{W}(J_2)$ or $(M, H, G) \in \mathcal{W}(J_3)$. Let us denote the class $\mathcal{W} := \bigcap_{\alpha=1}^3 \mathcal{W}(J_\alpha)$.

The next theorem gives a sufficient condition an almost (H, G) -manifold to be integrable.

Theorem 1.1 ⁽⁴⁾ *Let (M, H, G) belongs to the class $\mathcal{W}(J_\alpha) \cap \mathcal{W}(J_\beta)$. Then (M, H, G) is of class $\mathcal{W}(J_\gamma)$ for all cyclic permutations (α, β, γ) of $(1, 2, 3)$.*

Let us remark that necessary and sufficient conditions (M, H, G) to be in $\mathcal{W}$ are

$$\theta_\alpha \circ J_\alpha = -\frac{2n}{2n-1}\theta_1 \circ J_1, \quad \alpha = 2, 3. \quad (14)$$

2 Pseudo-hyper-Kähler manifolds

Definition 2.1 ⁽⁴⁾ *A pseudo-Hermitian manifold is called a pseudo-hyper-Kähler manifold, if $\nabla J_\alpha = 0$ ($\alpha = 1, 2, 3$) with respect to the Levi-Civita connection generated by g .*

It is clear, then $F_\alpha = 0$ ($\alpha = 1, 2, 3$) holds or the manifold is Kählerian with respect to J_α , i.e. $(M, H, G) \in \mathcal{K}(J_\alpha)$.

Immediately we obtain that if (M, H, G) belongs to $\mathcal{K}(J_\alpha) \cap \mathcal{W}(J_\beta)$ then $(M, H, G) \in \mathcal{K}(J_\gamma)$ for all cyclic permutations (α, β, γ) of $(1, 2, 3)$.

Then the following sufficient condition for a $\mathcal{K}$ -manifold is valid.

Theorem 2.1 ⁽⁴⁾ *If $(M, H, G) \in \mathcal{K}(J_\alpha) \cap \mathcal{W}(J_\beta)$ then M is a pseudo-hyper-Kähler manifold ($\alpha \neq \beta \in \{1, 2, 3\}$).*

Let (M^{4n}, H, G) be a pseudo-hyper-Kähler manifold and ∇ be the Levi-Civita connection generated by g . The curvature tensor seems as follows

$$R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]}Z, \quad (15)$$

and the corresponding tensor of type $(0, 4)$ is

$$R(X, Y, Z, W) = g(R(X, Y)Z, W), \quad \forall X, Y, Z, W \in \mathfrak{X}(M). \quad (16)$$

Lemma 2.2 *The curvature tensor of a pseudo-hyper-Kähler manifold has the following properties:*

$$\begin{aligned} R(X, Y, Z, W) &= R(X, Y, J_1 Z, J_1 W) = R(J_1 X, J_1 Y, Z, W) \\ &= -R(X, Y, J_2 Z, J_2 W) = -R(J_2 X, J_2 Y, Z, W) \\ &= -R(X, Y, J_3 Z, J_3 W) = -R(J_3 X, J_3 Y, Z, W), \end{aligned} \quad (17)$$

$$\begin{aligned} R(X, Y, Z, W) &= R(X, J_1 Y, J_1 Z, W) \\ &= -R(X, J_2 Y, J_2 Z, W) = -R(X, J_3 Y, J_3 Z, W). \end{aligned} \quad (18)$$

Proof. The equality (17) is valid, because of (15), (16), the condition $\nabla J_\alpha = 0$ ($\alpha = 1, 2, 3$), the equality (8) and the properties of the curvature $(0, 4)$ -tensor.

To prove (18), we will show at first that the property $R(X, J_2 Y, J_2 Z, W) = -R(X, Y, Z, W)$ holds. Indeed, from (17) we get

$$R(J_2 X, Y, Z, W) = R(X, J_2 Y, Z, W), \quad R(X, Y, J_2 Z, W) = R(X, Y, Z, J_2 W)$$

and $\mathfrak{S}_{X,Y,Z} R(X, Y, J_2 Z, J_2 W) = 0$, where $\mathfrak{S}_{X,Y,Z}$ denotes the cyclic sum regarding X, Y, Z . In the last equality we replace Y by $J_2 Y$ and W by $J_2 W$. We get

$$-R(X, J_2 Y, J_2 Z, W) - R(J_2 Y, Z, J_2 X, W) + R(Z, X, Y, W) = 0. \quad (19)$$

Replacing Y by Z , and inversely, we get

$$-R(X, J_2 Z, J_2 Y, W) - R(J_2 Z, Y, J_2 X, W) + R(Y, X, Z, W) = 0. \quad (20)$$

As we have

$$-R(J_2 Z, Y, J_2 X, W) = -R(Z, J_2 Y, J_2 X, W) = R(J_2 Y, Z, X, W),$$

with the help of (19) and (20) we obtain

$$\begin{aligned} &-R(X, J_2 Y, J_2 Z, W) - R(X, J_2 Z, J_2 Y, W) \\ &+ R(Z, X, Y, W) + R(Y, X, Z, W) = 0. \end{aligned} \quad (21)$$

According to the first Bianchi identity and (17), we obtain

$$\begin{aligned} -R(X, J_2 Z, J_2 Y, W) &= R(J_2 Z, J_2 Y, X, W) + R(J_2 Y, X, J_2 Z, W) \\ &= -R(Z, Y, X, W) - R(X, J_2 Y, J_2 Z, W). \end{aligned}$$

Then the equality (21) seem as follows

$$-2R(X, J_2 Y, J_2 Z, W) + R(Z, X, Y, W) - R(X, Y, Z, W) + R(Y, Z, X, W) = 0$$

By the first Bianchi identity the equality is transformed in the following

$$-2R(X, J_2 Y, J_2 Z, W) - 2R(X, Y, Z, W) = 0,$$

which is equivalent to

$$R(X, J_2Y, J_2Z, W) = -R(X, Y, Z, W). \quad (22)$$

As the tensor R has the same properties with respect to J_3 , and to J_2 , it follows that the next equality holds, too.

$$R(X, J_3Y, J_3Z, W) = -R(X, Y, Z, W). \quad (23)$$

Using (22) and (23) for $J_1 = J_2J_3$ we get successively that

$$\begin{aligned} R(X, Y, Z, W) &= R(X, J_1Y, J_1Z, W) \\ &= R(X, J_2(J_3Y), J_2(J_3Z), W) = -R(X, J_3Y, J_3Z, W), \end{aligned}$$

which completes the proof of (18).

Now we will prove a theorem which gives us a geometric characteristic of the pseudo-hyper-Kähler manifolds.

Theorem 2.3 *Each pseudo-hyper-Kähler manifold is a flat pseudo-Riemannian manifold with signature $(2n, 2n)$.*

Proof. Lemma 2.2 implies the properties

$$\begin{aligned} -R(X, Y, Z, W) &= R(X, J_1Y, Z, J_1W) \\ &= R(X, J_2Y, Z, J_2W) = R(X, J_3Y, Z, J_3W). \end{aligned} \quad (24)$$

As $J_1 = J_2J_3$, we also have the following

$$\begin{aligned} R(X, J_1Y, Z, J_1W) &= R(X, J_2(J_3Y), Z, J_2(J_3W)) \\ &= -R(X, J_3Y, Z, J_3W) = R(X, Y, Z, W). \end{aligned}$$

Comparing (24) with the last equality we receive

$$-R(X, Y, Z, W) = R(X, J_1Y, Z, J_1W) = R(X, Y, Z, W),$$

or $R \equiv 0$.

3 Conformal transformations of the pseudo-Hermitian metric

The usual conformal transformation $c : \bar{g} = e^{2u}g$, where u is a differential function on M^{4n} , is known. Since $g_\alpha(\cdot, \cdot) = g(J_\alpha \cdot, \cdot)$, the conformal transformation of g causes the same changes of the pseudo-Hermitian metrics g_2, g_3 and the Kähler form $\Phi \equiv g_1$. Then we say that it is given a conformal transformation c of G to $\bar{G}$ determined by $u \in \mathcal{F}(M)$. These conformal transformations form a group denoted by C . The hypercomplex pseudo-Hermitian manifolds (M, H, G) and $(M, H, \bar{G})$ we call C -equivalent manifolds or conformal-equivalent manifolds.

Let ∇ and $\bar{\nabla}$ be the Levi-Civita connections determined by the metrics g and $\bar{g}$, respectively. The known condition for a Levi-Civita connection implies the following relation

$$\bar{\nabla}_X Y = \nabla_X Y + du(X)Y + du(Y)X - g(X, Y)\text{grad}(u). \quad (25)$$

Using (25) and the definitions of structural tensors for ∇ and $\bar{\nabla}$ we obtain

$$\bar{F}_1(X, Y, Z) = e^{2u} [F_1(X, Y, Z) - g(X, Y)du(J_1 Z) + g(X, Z)du(J_1 Y) + g(J_1 X, Y)du(Z) - g(J_1 X, Z)du(Y)], \quad (26)$$

$$\bar{F}_\alpha(X, Y, Z) = e^{2u} [F_\alpha(X, Y, Z) + g(X, Y)du(J_\alpha Z) + g(X, Z)du(J_\alpha Y) - g(J_\alpha X, Y)du(Z) - g(J_\alpha X, Z)du(Y)] \quad (27)$$

for $\alpha = 2, 3$. The last two equalities imply the following relations for the corresponding structural 1-forms

$$\bar{\theta}_1 = \theta_1 - 2(2n-1)du \circ J_1, \quad \bar{\theta}_\alpha = \theta_\alpha + 4ndu \circ J_\alpha, \quad \alpha = 2, 3. \quad (28)$$

Let us denote the following (0,3)-tensors.

$$P_1(x, y, z) = F_1(x, y, z) - \frac{1}{2(2n-1)} [g(x, y)\theta_1(z) - g(x, z)\theta_1(y) - g(x, J_1 y)\theta_1(J_1 z) + g(x, J_1 z)\theta_1(J_1 y)], \quad (29)$$

$$P_\alpha(x, y, z) = F_\alpha(x, y, z) - \frac{1}{4n} [g(x, y)\theta_\alpha(z) + g(x, z)\theta_\alpha(y) + g(x, J_\alpha y)\theta_\alpha(J_\alpha z) + g(x, J_\alpha z)\theta_\alpha(J_\alpha y)], \quad \alpha = 2, 3. \quad (30)$$

According to (12) and (13) it is clear that

$$(M, H, G) \in \mathcal{W}(J_\alpha) \iff P_\alpha = 0 \quad (\alpha = 1, 2, 3).$$

The equalities (26)–(28) imply the following two interconnections

$$\bar{P}_\alpha = e^{2u} P_\alpha, \quad \alpha = 1, 2, 3; \quad (31)$$

$$\bar{\theta}_\alpha \circ J_\alpha + \frac{2n}{2n-1} \bar{\theta}_1 \circ J_1 = \theta_\alpha \circ J_\alpha + \frac{2n}{2n-1} \theta_1 \circ J_1, \quad \alpha = 2, 3. \quad (32)$$

From (31) we receive that each of $\mathcal{W}(J_\alpha)$ ($\alpha = 1, 2, 3$) is invariant with respect to the conformal transformations of C , i.e. they are C -invariant classes. Having in mind also (32), we state the validity of the following

Theorem 3.1 *The class $\mathcal{W}$ of hypercomplex pseudo-Hermitian manifolds is C -invariant.*

Now we will determine the class of the (locally) C -equivalent $\mathcal{K}$ -manifolds. Let us denote the following subclass $\mathcal{W}^0 := \{\mathcal{W} \mid d(\theta_1 \circ J_1) = 0\}$.

Theorem 3.2 *A hypercomplex pseudo-Hermitian manifold belongs to $\mathcal{W}^0$ if and only if it is C -equivalent to a pseudo-hyper-Kähler manifold.*

Proof. Let (M, H, G) be a pseudo-hyper-Kähler manifold, i.e. $(M, H, G) \in \mathcal{K}$. Then $F_\alpha = \theta_\alpha = 0$ ($\alpha = 1, 2, 3$). Hence (28) has the form

$$\bar{\theta}_1 = -2(2n-1)du \circ J_1, \quad \bar{\theta}_\alpha = 4ndu \circ J_\alpha, \quad \alpha = 2, 3. \quad (33)$$

From (26), (27) and (33) and having in mind (12) and (13) we obtain that $(M, H, \bar{G})$ is a $\mathcal{W}$ -manifold. According to (33) the 1-forms $\bar{\theta}_\alpha \circ J_\alpha$ ($\alpha = 1, 2, 3$) are closed. Because of (14) the condition $d(\bar{\theta}_1 \circ J_1) = 0$ is sufficient.

Conversely, let $(M, H, \bar{G})$ be a $\mathcal{W}$ -manifold with closed $\bar{\theta}_1 \circ J_1$. Because of (14) the 1-forms $\bar{\theta}_\alpha \circ J_\alpha$ ($\alpha = 2, 3$) are closed, too. We determine the function u as a solution of the differential equation $du = -\frac{1}{2(2n-1)}\bar{\theta}_1 \circ J_1$. Then by an immediate verification we state that the transformation $c^{-1} : g = e^{-2u}\bar{g}$ converts $(M, H, \bar{G})$ into $(M, H, G) \in \mathcal{K}$. This completes the proof.

Let us remark the following inclusions

$$\mathcal{K} \subset \mathcal{W}^0 \subset \mathcal{W} \subset \mathcal{W}(J_\alpha), \quad \alpha = 1, 2, 3.$$

Let R, ρ, τ and $\bar{R}, \bar{\rho}, \bar{\tau}$ be the curvature tensors, the Ricci tensors, the scalar curvatures corresponding to ∇ and $\bar{\nabla}$, respectively. The following tensor is curvature-like, i.e. it has the same properties as R .

$$\begin{aligned} \psi_1(S)(X, Y, Z, U) &= g(Y, Z)S(X, U) - g(X, Z)S(Y, U) \\ &\quad + g(X, U)S(Y, Z) - g(Y, U)S(X, Z), \end{aligned}$$

where S is a symmetric tensor.

Having in mind (25) and (15), we obtain

Proposition 3.3 *The following relations hold for the C -equivalent (H, G) -manifolds*

$$\begin{aligned} \bar{R} &= e^{2u}\{R - \psi_1(S)\}, \\ \bar{\rho} &= \rho - \text{tr}Sg - 2(2n-1)S, \quad \bar{\tau} = e^{-2u}\{\tau - 2(4n-1)\text{tr}S\}, \end{aligned} \quad (34)$$

where

$$S(Y, Z) = S(Z, Y) = (\nabla_Y du)Z + du(Y)du(Z) - \frac{1}{2}du(\text{grad}(du))g(Y, Z). \quad (35)$$

If (M, H, G) is a C -equivalent $\mathcal{W}$ -manifold to a $\mathcal{K}$ -manifold, i.e. $(M, H, G) \in \mathcal{W}^0$, then Proposition 3.3 implies

Corollary 3.4 *A $\mathcal{W}^0$ -manifold has the following curvature characteristic*

$$R = \frac{1}{2(2n-1)} \left\{ \psi_1(\rho) - \frac{\tau}{4n-1} \pi_1 \right\},$$

where $\pi_1(X, Y, Z, U) = \frac{1}{2}\psi_1(g) = g(Y, Z)g(X, U) - g(X, Z)g(Y, U)$.

It is well known that the C -invariant tensor of each pseudo-Riemannian manifold is the so-called Weil tensor W . From (34) we receive immediately

$$\bar{W} = e^{2u}W, \quad W = R - \frac{1}{2(2n-1)} \left\{ \psi_1(\rho) - \frac{\tau}{4n-1} \pi_1 \right\}. \quad (36)$$

Let us remark that the vanishing of the Weil tensor is a necessary and sufficient condition a pseudo-Riemannian manifold to be conformal equivalent to a flat manifold with dimension greater than 3.

This is confirmed by the combining of Theorem 2.3, Theorem 3.2 and Corollary 3.4, i.e. $(M, H, G) \in \mathcal{W}^0$ iff $W = 0$ on (M, H, G) .

Since each conformal transformation determines uniquely a symmetric tensor S by (35) then it takes an interest in the consideration S as a bilinear form on $T_p M$ belonging to each of the components L_α , $(\alpha = 0, 1, 2, 3)$.

Let $S \in L_0$. In view of (5) $\text{tr} S = 0$ holds and according to (34) we receive $\bar{\tau} = e^{-2u}\tau$ and an invariant tensor $W_0 = R - \frac{1}{2(2n-1)}\psi_1(\rho)$. When W_0 vanishes on (M, H, G) then the curvature tensor has the form $R = \frac{1}{2(2n-1)}\psi_1(\rho)$.

In the cases when $S \in L_\alpha$ ($\alpha = 1, 2, 3$) we consider (M, H, G) as an $\mathcal{W}^0$ -manifold. Then according to Theorem 2.3 and Theorem 3.2 we have $\bar{R} = 0$ on the C -equivalent $\mathcal{K}$ -manifold of (M, H, G) .

Now let $S \in L_1$. By reason of $g \in L_1$ we have a cause for the consideration of the possibility $S = \lambda g$. Hence $\lambda = \frac{\text{tr} S}{4n} = \frac{\tau}{8n(4n-1)}$. Then having in mind (34) $R = \frac{\tau}{4n(4n-1)}\pi_1$ holds true. From here it is clear that if $S \in L_1$ then (M, H, G) is an Einstein manifold.

Let us consider the case when $S \in L_2$. Then according to (6) $\text{tr} S$ vanishes, and from (34) τ vanishes, too. Because of $g_3 \in L_2$ we consider $S = \lambda g_3$, whence $\lambda = -\frac{\text{tr}(S \circ J_3)}{4n}$. Then (34) implies $R = \frac{\text{tr}(S \circ J_3)}{4n}\pi_3^{J_3}$, where $\pi_3^{J_3}$ is the following tensor π_3 with respect to the complex structure $J = J_3$

$$\pi_3(X, Y, Z, U) = -\pi_1(X, Y, JZ, U) - \pi_1(X, Y, Z, JU).$$

It is known² that π_3 is a Kähler curvature-like tensor, i.e. it satisfies the property $\pi_3(X, Y, JZ, JU) = -\pi_3(X, Y, Z, U)$. Therefore in this case R is Kählerian with respect to J_3 and the tensor $R^{*J_3} : R^{*J_3}(X, Y, Z, U) = R(X, Y, Z, J_3 U)$ is curvature-like. Then we obtain immediately

$$R = \frac{\tau(R^{*J_3})}{8n(2n-1)}\pi_3^{J_3}, \quad \rho = -\frac{\tau(R^{*J_3})}{4n}g_3.$$

Hence if $S \in L_2$ then (M, H, G) is a $*$ -Einstein manifold with respect to J_3 .

By an analogous way, in the case when $S \in L_3$ we receive that (M, H, G) is a $*$ -Einstein manifold with respect to J_2 .

4 A 4-dimensional pseudo-Riemannian spherical manifold with (H, G) -structure

In $\mathbb{R}^5$ is considered a hypersurface S_2^4 in $\mathbb{R}_2^5$ by the equation

$$-(z^1)^2 - (z^2)^2 + (z^3)^2 + (z^4)^2 + (z^5)^2 = 1, \quad (37)$$

where $Z(z^1, z^2, z^3, z^4, z^5)$ is the positional vector of $p \in S_2^4$.

Let (u^1, u^2, u^3, u^4) be local coordinates of p on S_2^4 . The hypersurface S_2^4 is defined by the scalar parametric equations:

$$\begin{aligned} z^1 &= \sinh u^1 \cos u^2, & z^2 &= \sinh u^1 \sin u^2, & z^3 &= \cosh u^1 \cos u^3 \cos u^4, \\ z^4 &= \cosh u^1 \cos u^3 \sin u^4, & z^5 &= \cosh u^1 \sin u^3. \end{aligned} \quad (38)$$

Further we consider the manifold on $\tilde{S}_2^4 = S_2^4 \setminus \{(0, 0, 0, 0, \pm 1)\}$, i.e. we omit two points for which $\{u^1 \neq 0\} \cap \{u^3 \neq (2k+1)\pi/2, k \in \mathbb{Z}\}$. The tangent space $T_p \tilde{S}_2^4$ of $\tilde{S}_2^4$ in the point $p \in \tilde{S}_2^4$ is determined by the vectors $z_i = \frac{\partial Z}{\partial u^i} (i = 1, 2, 3, 4)$. The vectors z_i are linearly independent on $\tilde{S}_2^4$, defined by (38), and $T_p \tilde{S}_2^4$ has a basis (z_1, z_2, z_3, z_4) in every point $p \in \tilde{S}_2^4$.

The restriction of $\langle \cdot, \cdot \rangle$ from $\mathbb{R}_2^5$ to S_2^4 is a pseudo-Riemannian metric g on S_2^4 with signature $(2, 2)$. The non-zero components $g_{ij} = \langle z_i, z_j \rangle$ are

$$g_{11} = -1, \quad g_{22} = -\sinh^2 u^1, \quad g_{33} = \cosh^2 u^1, \quad g_{44} = \cosh^2 u^1 \cos^2 u^3. \quad (39)$$

The hypersurface S_2^4 is equipped with an almost hypercomplex structure $H = (J_\alpha), (\alpha = 1, 2, 3)$, where the non-zero components of the matrix of J_α with respect to the local basis $\{\frac{\partial}{\partial u^i}\}_{i=1}^4$ are

$$\begin{aligned} (J_1)_2^1 &= -\frac{1}{(J_1)_1^2} = -\sinh u^1, & (J_1)_4^3 &= -\frac{1}{(J_1)_3^4} = \cos u^3, \\ (J_2)_3^1 &= -\frac{1}{(J_2)_1^3} = -\cosh u^1, & (J_2)_4^2 &= -\frac{1}{(J_2)_2^4} = -\coth u^1 \cos u^3, \\ (J_3)_4^1 &= -\frac{1}{(J_3)_1^4} = \cosh u^1 \cos u^3, & (J_3)_2^3 &= -\frac{1}{(J_3)_3^2} = \tanh u^1. \end{aligned} \quad (40)$$

Theorem 4.1 ⁽⁴⁾ *The spherical pseudo-Riemannian 4-dimensional manifold, defined by (38), admits a hypercomplex pseudo-Hermitian structure on $\tilde{S}_2^4$, determined by (40) and (39), with respect to which it is of the class $\mathcal{W}(J_1)$ but it does not belong to $\mathcal{W}$ and it has a constant sectional curvature $k = 1$.*

Let us consider a conformal transformation determined by the function u which is a solution of the equation $du = -\frac{1}{2(2n-1)}(\theta_1 \circ J_1)$, where the nonzero component of θ_1 with respect to the local basis $\{\frac{\partial}{\partial u^i}\}$ ($i = 1, 2, 3, 4$) is $\theta_1(\frac{\partial}{\partial u^2}) = \frac{2 \sinh^2 u^1}{\cosh u^1}$.

Since $\tilde{S}_2^4$ has a constant sectional curvature then the Weil tensor is vanishes, i.e. $\tilde{S}_2^4$ is C -equivalent to a flat $\mathcal{K}(J_1)$ -manifold. If we admit that it is in

$\mathcal{K}$, then according to Theorem 3.2 we obtain that the manifold $(\tilde{S}_2^4, H, G) \in \mathcal{W}$ which is a contradiction. Therefore the considered manifold is C -equivalent to a flat $\mathcal{K}(J_1)$ -manifold, but it is not a pseudo-hyper-Kähler manifold. By direct verification we state that the tensor S of this conformal transformation belongs to L_1 . Therefore $(\tilde{S}_2^4, H, G)$ is an Einstein manifold.

References

1. D. V. Alekseevsky and S. Marchiafava, *Quaternionic structures on a manifold and subordinated structures*, Ann. Mat. Pura Appl. (IV), CLXXI (1996), 205–273.
2. G. Ganchev and A. Borisov, *Note on the almost complex manifolds with a Norden metric*, Compt. rend. Acad. bulg. Sci., **39** (1986), no. 5, 31–34.
3. A. Gray and L. M. Hervella, *The sixteen classes of almost Hermitian manifolds and their linear invariants*, Ann. Mat. Pura Appl. (IV), CXXIII (1980), 35–58.
4. K. Gribachev, M. Manev and S. Dimiev, *Almost hypercomplex pseudo-Hermitian manifolds*, Boll. Unione Mat. Ital. Sez. A, (to appear)
5. A. Sommese, *Quaternionic manifolds*, Math. Ann. **212** (1975), 191–214.
6. J. Wolf, *Spaces of constant curvature*, University of California, Berkley, California, 1972.